Sanitized - Approved For Release: CIA-KDI-83-00423R00 500550018 3/10 TUDENTIAL CIA 25X1X6

Country: Brazil

Subject: Dr Repul Tutrut, Natal State Mineral Prospector/

Possible Bauxite Deposits in Rio Grande do Norte

DATE 2 Jul 54 8 Jul 54

Place Acquired:

Date Acquired :

Date of Info : May 54

A#45247

25X1X6

SOURCE: US mining engineer.

<u> 25X1A</u>

25X1X6

25X1X

25X1X

ı.

is employed within the Department

of Agriculture of Rio Grande do Norte as a practical mine He is a graduate of the Institute Schneider in Paris, which was founded by the French industrialist Schneider Caruso. He has worked on mineral prospecting in many parts of French Africa, Madagascar, French Guiana and Surinam. He has never worked on bauxite specifically but has observed bauxite in place in Surinem.

- 25X1A is a little slow in getting around on foot due to a war injury, but he has covered most of the state by motorcycle. All of the reports of bauxite in Rio Grande do Norte have originated with him. These reports are based on visual recognition only and are not backed up by any analyses. The state does not have a chemical laboratory, although he personally has a mineral testing kit. Tutrut rates the supposed bauxite occurrences in the following order of importance.
 - e. Itaretama
 - b. Açu
 - e. Júcurutu
 - d. Nisea Floresta
- 3. I drove around the country a little and examined a few laterites exposed in highway cuts and borrow pits. The GSA map of South America maps the coastal fringe in this area as "undifferentiated tertiary." What I saw was an alteration of almost pure quartz sands. Clays and laterities at the surface are just under a shallow soil cover. In many places dune sand from the beaches has blown in to add some extra quartz sand to the laterite. These laterites, which are hard enough to be used for road surfacing, are not of the red

JUL 23 1954

Sanitized - Approved For Release : CIA-RDP83-00423R001500560011-2

variety but are more cream or tan in color. I did not observe them to be pisolitic in any exposure. Only somewhat extensive sampling could disclose their true nature. The topography is favorable - flattopped hills 50-100 feet above sea level, dissected by widely spaced, small streams that cut almost to sea level. A great deal of this area is cultivated with corn, manioc, etc. There are a few carnauba wax pods. The rainfall is sufficient for crops, but there is no dense jungle.

-end-

Sulip Gen'l Phining Mgo -40 - 45,1 Orea VYFE ACT-DATE INFO-TFR-BATE AREA

Bray i), Rio Grande - 5-4 - 5-4

036 - 39.1

-2-